



## Corrigendum

Corrigendum to “Onset of  $\eta$ -nuclear binding in a pionless EFT approach” [Phys. Lett. B 771 (2017) 297–302]N. Barnea<sup>a</sup>, B. Bazak<sup>b</sup>, E. Friedman<sup>a</sup>, A. Gal<sup>a,\*</sup><sup>a</sup> Racah Institute of Physics, The Hebrew University, 91904 Jerusalem, Israel<sup>b</sup> IPNO, CNRS/IN2P3, Univ. Paris-Sud, Université Paris-Saclay, F-91406 Orsay, France

## ARTICLE INFO

## Article history:

Received 19 September 2017

Accepted 3 October 2017

Available online 19 October 2017

Editor: W. Haxton

## Keywords:

Few-body systems

Mesic nuclei

EFT calculations

## ABSTRACT

A three-body force acting between the  $\eta$ -meson and two nucleons was overlooked inadvertently in the model description and discussion in the published version of our paper “Onset of  $\eta$ -nuclear binding in a pionless EFT approach” [Phys. Lett. B 771 (2017) 297–302] while present in the actual numerical calculations. The stated conclusion that a stabilizing  $\eta NN$  contact term was not needed is therefore incorrect. Such a three-body force, associated with a new low energy constant  $d_{\eta NN}^\Lambda$ , must be introduced at leading order to stabilize  $\eta$ -nucleus systems.


© 2017 The Author(s). Published by Elsevier B.V. All rights reserved.

A three-body  $\eta NN$  force was inadvertently overlooked in the potential model description and discussion in Ref. [1]. In the actual calculations, however, the leading order interaction between the  $\eta$  and the nucleons was composed of the  $\eta N$  term discussed in Sect. 2.3, supplemented by an  $\eta NN$  term

$$V_{\eta N_i N_j} = d_{\eta NN}^\Lambda \delta_\Lambda(r_{\eta N_i}, r_{\eta N_j}). \quad (1)$$

Here,  $\delta_\Lambda(r_{\eta N_i}, r_{\eta N_j})$  is a product of normalized pairwise Gaussians  $\delta_\Lambda(r_{\eta N_i})$  and  $\delta_\Lambda(r_{\eta N_j})$ , with range parameter inversely proportional to the momentum-scale parameter  $\Lambda$ , as defined by Eq. (4) of Ref. [1]. For the results presented in the paper, the low energy constant (LEC)  $d_{\eta NN}^\Lambda$  was set equal to the nuclear  $NNN$  LEC  $d_3^\Lambda$ . Setting  $d_{\eta NN}^\Lambda = 0$ , the  $\eta$ -deuteron ( $\eta d$ ) system, and therefore any  $\eta$ -nucleus system, would collapse as  $\Lambda \rightarrow \infty$ .

The parameter  $d_{\eta NN}^\Lambda$  is a free parameter to be fixed by experimental data. In the absence of such data one may estimate its value using the nuclear  $NNN$  LEC,  $d_{\eta NN}^\Lambda = d_3^\Lambda$ , as done in [1], or to set a bound on its value accepting that  $\eta d$  is unbound [2], i.e. set  $d_{\eta NN}^\Lambda$  so that  $B_\eta(\eta d) = 0$ . To check the sensitivity of the results in [1] to these distinct choices of  $d_{\eta NN}^\Lambda$ , we present in Figs. 1 and 2 calculations of  $\eta$  separation energies  $B_\eta$  in  $\eta^3\text{He}$  and  $\eta^4\text{He}$ ,



**Fig. 1.**  $B_\eta(\eta^3\text{He})$  as a function of  $1/\Lambda$ , calculated using  $\eta N$  potentials  $v_{\eta N}^{\text{GW}}(E)$  for two choices of the  $\eta NN$  LEC. Solid lines:  $d_{\eta NN}^\Lambda = d_3^\Lambda$  [1], dashed lines:  $d_{\eta NN}^\Lambda$  fitted to produce  $B_\eta(\eta d) = 0$ . Self consistent calculations are marked by squares (red); calculations using threshold values  $v_{\eta N}^{\text{GW}}(E_{\text{th}})$  are marked by spheres (blue). Linear extrapolations to a point-like interaction,  $\Lambda \rightarrow \infty$ , are marked by straight lines.

respectively, using  $\eta N$  potentials  $v_{\eta N}^{\text{GW}}(E)$  under these two choices of  $d_{\eta NN}^\Lambda$ . Figs. 1 and 2 update the original Figs. 4 and 5 in [1].

Figs. 1 and 2 demonstrate that the two choices made for the three-body  $\eta NN$  LEC yield practically identical values of  $B_\eta$  in the limit  $\Lambda \rightarrow \infty$ . For values of  $\Lambda$  near the physical breakdown scale  $\Lambda \approx 4 \text{ fm}^{-1}$ , however,  $B_\eta$  differs by about 0.7 MeV for  $\eta^3\text{He}$  and 2 MeV for  $\eta^4\text{He}$  between the two choices applied in self consistent

DOI of original article: <https://doi.org/10.1016/j.physletb.2017.05.066>.

\* Corresponding author.

E-mail address: [avragal@savion.huji.ac.il](mailto:avragal@savion.huji.ac.il) (A. Gal).




Fig. 2. Same as in Fig. 1, but for  $\eta^4\text{He}$  instead of  $\eta^3\text{He}$ .

calculations (lower group of curves). Since  $\eta d$  is unbound [2], the choice marked in dashed lines in both figures is likely to somewhat overestimate  $B_\eta$ . Nevertheless, these  $\eta$  separation energies are in good agreement with the non-EFT  $B_\eta$  values calculated recently using the same two-body energy dependent  $\eta N$  interaction [3].

**References**

[1] N. Barnea, B. Bazak, E. Friedman, A. Gal, Phys. Lett. B 771 (2017) 297.  
 [2] B. Krusche, C. Wilkin, Prog. Part. Nucl. Phys. 80 (2015) 43.  
 [3] N. Barnea, E. Friedman, A. Gal, Nucl. Phys. A 968 (2017) 35.